

Anti-ETFA antibody [2B11AE8] ab110316

敲除验证

8 References 5 图像

概述	
产品名称	Anti-ETFA抗体[2B11AE8]
描述	小鼠单克隆抗体[2B11AE8] to ETFA
宿主	Mouse
经测试应用	适用于: WB, ICC/IF, Flow Cyt, IHC-P
种属反应性	与反应: Mouse, Rat, Cow, Human
免疫原	Recombinant full length protein. This information is considered to be commercially sensitive.
阳性对照	ICC: HeLa cells. Flow Cyt: HL-60 cells. IHC-P: Human heart tissue. WB: HepG2 and HEK-293T whole cell lysates; Human liver mitochondria; Bovine heart mitochondria; Rat liver mitochondria; Mouse liver mitochondria.
常规说明	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> <p>Product was previously marketed under the MitoSciences sub-brand.</p>
性能	
形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.5 Preservative: 0.02% Sodium azide Constituent: HEPES buffered saline
纯度	Proprietary Purification

纯化说明	The antibody was produced in vitro using hybridomas grown in serum-free medium, and then purified by biochemical fractionation. Purity >95% by SDS-PAGE.
克隆	单克隆
克隆编号	2B11AE8
同种型	IgG2b
轻链类型	kappa

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab110316于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 35 kDa.
ICC/IF		Use a concentration of 4 µg/ml.
Flow Cyt		Use a concentration of 1 µg/ml. ab170192 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.
IHC-P		Use a concentration of 5 µg/ml.

靶标

功能	The electron transfer flavoprotein serves as a specific electron acceptor for several dehydrogenases, including five acyl-CoA dehydrogenases, glutaryl-CoA and sarcosine dehydrogenase. It transfers the electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase).
疾病相关	Glutaric aciduria 2A
序列相似性	Belongs to the ETF alpha-subunit/FixB family.
结构域	Domain I shares an identical polypeptide fold with the beta subunit ETFB though there is no sequence similarity.
翻译后修饰	The N-terminus is blocked.
细胞定位	Mitochondrion matrix.

图片



Western blot - Anti-ETFA antibody [2B11AE8] (ab110316)

All lanes : Anti-ETFA antibody [2B11AE8] (ab110316) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 2 : ETFA knockout HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lysates/proteins at 40 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed ([ab216777](#)) at 1/10000 dilution

Predicted band size: 35 kDa

Observed band size: 35 kDa

Lanes 1-2: Merged signal (red and green). Green - ab110316 observed at 35 kDa. Red - loading control [ab52901](#) observed at kDa.

ab110316 Anti-ETFA antibody [2B11AE8] was shown to specifically react with ETFA in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line [ab266513](#) (knockout cell lysate [ab257943](#)) was used. Wild-type and ETFA knockout samples were subjected to SDS-PAGE. ab110316 and Anti-beta Tubulin [EP1331Y] - Microtubule Marker ([ab52901](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed ([ab216777](#)) and Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed ([ab216772](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-ETFA antibody [2B11AE8] (ab110316)

All lanes : Anti-ETFA antibody [2B11AE8] (ab110316) at 1 µg/ml

Lane 1 : Human heart tissue

Lane 2 : HepG2 whole cells

Lane 3 : Human liver mitochondria


Lane 4 : Bovine heart mitochondria

Lane 5 : Rat liver mitochondria

Lane 6 : Mouse liver mitochondria


Lysates/proteins at 10 µg per lane.

Predicted band size: 35 kDa




Immunocytochemistry/ Immunofluorescence - Anti-ETFA antibody [2B11AE8] (ab110316)

Immunocytochemistry image of stained HeLa cells. The cells were paraformaldehyde fixed (4%, 20 minutes) and Triton X-100 permeabilized (0.1%, 15 minutes). The cells were incubated with the antibody (ab110316, 4 µg/mL) for 2 hours at room temperature or over night at 4°C. The secondary antibody was (red) Alexa Fluor® 594 goat anti-mouse IgG (H+L) at a 1/1000 dilution for 1 hour. 10% Goat serum was used as the blocking agent for all blocking steps. The target protein locates to the mitochondrial matrix



Flow Cytometry - Anti-ETFA antibody [2B11AE8] (ab110316)

HL-60 cells were stained with 1 µg/mL ab110316 (blue) or an equal amount of an isotype control antibody (red) and analyzed by flow cytometry.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ETFA antibody [2B11AE8] (ab110316)

IHC image of ETFA staining in Human heart formalin fixed paraffin embedded tissue section, performed on a Leica BondTM system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab110316, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors